

CHALMERS, GÖTEBORGS UNIVERSITET

EXAM for DYNAMICAL SYSTEMS

COURSE CODES: **TIF 155, FIM770GU, PhD**

| | |
|--------------------------|---|
| Time: | April 06, 2018, at 14 ⁰⁰ – 18 ⁰⁰ |
| Place: | Johanneberg |
| Teachers: | Kristian Gustafsson, 070-050 2211 (mobile), visits once around 15 ⁰⁰ |
| Allowed material: | Mathematics Handbook for Science and Engineering |
| Not allowed: | any other written material, calculator |

Maximum score on this exam: 12 points (need 5 points to pass).

Maximum score for homework problems: 24 points (need 10 points to pass).

CTH ≥ 18 passed; ≥ 26 grade 4; ≥ 31 grade 5,


GU ≥ 18 grade G; ≥ 28 grade VG.

1. Multiple choice questions [2 points] For each of the following questions identify **all** the correct alternatives A–E. Answer with letters among A–E. Some questions may have **more than one correct alternative**. In these cases answer with all appropriate letters among A–E.

a) Classify the fixed point of the three-dimensional dynamical system:

$$\dot{\mathbf{x}} = \mathbb{A}\mathbf{x}, \quad \text{where } \mathbb{A} = \begin{pmatrix} 1 & 0 & -2 \\ 0 & -1 & 0 \\ 2 & 0 & -3 \end{pmatrix}.$$


- A. It is a saddle point (all eigenvalues are real but of different signs).
 - B. It is a stable spiral (eigenvalues form a complex pair and the real parts of all eigenvalues are negative).
 - C. It is an unstable spiral (eigenvalues form a complex pair and the real parts of all eigenvalues are positive).
 - D. It is a stable node (all eigenvalues are negative).
 - E. It is an unstable node (all eigenvalues are positive).
- b) The figure below shows a bifurcation diagram for a dynamics in x with a parameter r . Solid lines show stable fixed points, dashed lines show unstable fixed points.



Consider the path (arrows) obtained by increasing r above r_s and then decreasing r again. Where does the system end up when r becomes smaller than zero?

- A. The system ends up at either the lower or upper branches of fixed points with 50% probability for each
- B. The system ends up either over or under the upper branch of fixed points with 50% probability for each
- C. The system ends up over the upper branch of fixed points
- D. The system ends up under the upper branch of fixed points
- E. The system ends up at the middle branch of fixed points

c) The figure below shows a flow with a single fixed point at its center:



What is the index of this fixed point if the trajectories are traversed backwards in time?

- A. -2
- B. -1
- C. 0
- D. 1
- E. 2

d) Consider the two-dimensional system

$$\dot{x} = y, \quad \dot{y} = x + x^4.$$

Which of the following is a conserved quantity of this system?

- A. $y + x + x^4$
- B. $y - x - x^4$
- C. $y^2/2 + x + x^4$
- D. $y^2/2 + 1 + 4x$
- E. $y^2/2 - x^2/2 - x^5/5$

- e) A three-dimensional dynamical system has the following Lyapunov exponents: $\lambda_1 = 0$, $\lambda_2 = -1$, $\lambda_3 = -2$. Which of the following statements are true?
- A. The system may be chaotic.
 - B. The system may be volume preserving.
 - C. The system may be Hamiltonian.
 - D. The system may have a globally attracting limit cycle.
 - E. The Lyapunov spectrum is unchanged if time changes sign.
- f) Which of the following statements about the generalized dimension spectrum D_q are true?
- A. For a multifractal, D_q increases with increasing q
 - B. For a monofractal, D_q is independent of q
 - C. When $q > 0$ low-density regions of the attractor gives the dominant contribution to D_q
 - D. When $q > 0$ high-density regions of the attractor gives the dominant contribution to D_q
 - E. The Kaplan-Yorke conjecture states that, in most cases, the correlation dimension is equal to D_1 .

2. Short questions [2 points] For each of the following questions give a concise answer within a few lines per question.

- a) Explain a method that can be used to decrease the dimensionality of a dynamical system. What assumptions does your method rely on?
- b) Explain what it means that a dynamical system shows hysteresis. Give an example of a system with hysteresis.
- c) What are the stable manifolds of a fixed point?
- d) Give two examples of different types of global bifurcations. Explain how one can experimentally distinguish the different bifurcations.
- e) Explain what a fractal (strange) attractor is and how it may form in a dynamical system.
- f) In a dynamical system of dimensionality larger than two, what does the **second** largest Lyapunov exponent characterize?

3. Bifurcations [2 points] Consider the system

$$\dot{r} = \mu r - gr^3 - r^5. \quad (1)$$

Assume that $r \geq 0$, and that μ and g are real parameters.

- Sketch the bifurcation diagram for the system (1) for the two cases $g = +1$ and $g = -1$. Identify the types of all bifurcations that occur in your bifurcation diagrams.
- When $g = 0$ a special kind of bifurcation occurs. For the two cases of $g = 0$ and a finite value $g > 0$, analytically determine how the locations of the fixed points depend on μ close to the bifurcation at $\mu = 0$. Sketch the bifurcation diagrams for the cases of $g = 0$ and a finite value $g > 0$ in the same plot.
- Discuss what would happen if a small imperfection h is added to the system (1) for the case $g = 0$:

$$\dot{r} = h + \mu r - r^5.$$

Make qualitative sketches for the cases $h > 0$ and $h < 0$.

4. Model of a national economy [2.5 points] A simple model for a national economy is provided by

$$\begin{aligned} \dot{I} &= I - \alpha C \\ \dot{C} &= \beta(I - C - G(I)), \end{aligned} \quad (2)$$

where $I \geq 0$ is the national income, $C \geq 0$ is the rate of consumer spending, and $G(I) \geq 0$ is the rate of government spending. Assume that $\alpha > 1$ and $\beta \geq 1$.

- Show that if $G(I) = G_0 = \text{const.}$, the system (2) has a single fixed point. Classify this fixed point in terms of α and β . Make a diagram in α - β -space that shows the regions of the different types of fixed points you find.
- Now consider the case of a government spending that increases linearly with the national income: $G(I) = G_0 + kI$, where $k > 0$. Show that there exists a value k_c such that for $k > k_c$ there exists no positive fixed points in the system. Does the system have any attractors when $k > k_c$?
- Solve the dynamics in subtask b) for $I(t)$ and $C(t)$ as functions of time in terms of a matrix exponential. Explicitly write out the components of the solution for the case $k = 1$, $\alpha = 2$ and $\beta = 1$. What is the long-term fate of the dynamics according to this solution? Is the result consistent with your result in subtask b)?

5. Damped driven pendulum [1.5 points] The equation of motion for the angle θ of a driven pendulum may be approximated by

$$\ddot{\theta} = -\frac{a}{m}\dot{\theta} - \frac{g}{b}\sin\theta + \frac{c}{I_0}. \quad (3)$$

Assume that m is the mass of the pendulum, g is the gravitational acceleration and I_0 is the moment of inertia of the pendulum.


- a) Briefly explain the origin of the three terms on the right hand side. Determine the dimensions of the parameters a , b , and c and give plausible physical interpretations of these three parameters.
- b) Show that Eq. (3) can be reduced to a dimensionless two-dimensional flow

$$\begin{aligned} \frac{d\theta}{dt'} &= y \\ \frac{dy}{dt'} &= -\sin\theta - \alpha y + I \end{aligned}$$

by a proper rescaling $t = t_0 t'$ and by suitable choices of the time-dependent function $y(t')$ and of the dimensionless parameters α and I . Explicitly check that α and I are dimensionless.

- c) Explain how this system may undergo a homoclinic bifurcation if the pendulum is weakly driven (consider fixed points separated by 2π in the θ -direction to be the same fixed point). Explain a method that can be used to find the corresponding bifurcation point. An explanation is enough, you do not need to explicitly evaluate the bifurcation point.

6. Koch curve [2 points] The figure below shows the first few generations in the construction of the Koch curve. It is obtained by, at each generation, replacing the central 1/3 of each line with two new lines joined at a bend angle $\alpha = \pi/3$. The fractal set is obtained by iterating to generation S_n with $n \rightarrow \infty$.



- Analytically find the box-counting dimension D_0 for the Koch curve.
- Now consider a general bend angle α . Assume that all lines in a given generation have the same lengths. Analytically, find the box-counting dimension as a function of the bend angle.
- Evaluate the box-counting dimension you obtained in subtask b) for the special cases $\alpha = 0$, $\alpha = \pi/3$ and $\alpha = \pi$. Discuss the results.
- Analytically find the area between the Koch curve and the baseline (the line in S_0) as a function of the bend angle α . You can assume that the length of the baseline is $L_0 = 1$.